arXiv:1606.07201v1 [math.RA] 23 Jun 2016

Hyperinvariant, characteristic and marked subspaces

Pudji Astuti
Faculty of Mathematics
and Natural Sciences
Institut Teknologi Bandung
Bandung 40132
Indonesia

Harald K. Wimmer
Mathematisches Institut
Universität Würzburg
D-97074 Würzburg
Germany

June 24, 2016

Abstract

Mathematical Subject Classifications (2000): 15A18, 47A15 15A57. 13C99

Keywords: hyperinvariant subspaces, marked subspaces, characteristic subspaces, invariant subspaces, Jordan basis.

Running title: Hyperinvariant subspaces

Abstract: Let V be a finite dimensional vector space over a field K and f a K-endomorphism of V. In this paper we study three types of f-invariant subspaces, namely hyperinvariant subspaces, which are invariant under all endomorphisms of V that commute with f, characteristic subspaces, which remain fixed under all automorphisms of V that commute with f, and marked subspaces, which have a Jordan basis (with respect to $f_{|X}$) that can be extended to a Jordan basis of V. We show that a subspace is hyperinvariant if and only if it is characteristic and marked. If K has more than two elements then each characteristic subspace is hyperinvariant.

1 Introduction

Let V be an n-dimensional vector space over a field K and let $f : V \to V$ be K-linear. We assume that the characteristic polynomial of f splits over K such that all eigenvalues of f are in K. In this paper we deal with three types of f-invariant subspaces, namely with hyperinvariant, characteristic and marked subspaces. To describe these three concepts we use the following notation. Let $\mathrm{Inv}(V)$ be the lattice of f-invariant subspaces of V and let $\mathrm{End}_f(V)$ be the algebra of all endomorphisms of V that commute with f. If a subspace X remains invariant for all $g \in \mathrm{End}_f(V)$ then X is called *hyperinvariant* for f [13, p. 305]. Let Hinv(V) be the set of hyperinvariant subspaces of V. It is obvious that Hinv(V) is a lattice. Because of $f \in \mathrm{End}_f(V)$ we have Hinv(V) $\subseteq$ Inv(V). We refer to [13], [9], [17], [19] for results on hyperinvariant subspaces. The group of automorphisms of V that commute with f will be denoted by $\mathrm{Aut}_f(V)$. A subspace X of V will be called *characteristic* (with respect to f) if $X \in \mathrm{Inv}(V)$ and $\alpha(X) = X$ for all $\alpha \in \mathrm{Aut}_f(V)$. Let Chinv(V) be set of characteristic subspaces of V. Obviously, also Chinv(V) is a lattice, and Hinv(V) $\subseteq$ Chinv(V).

Set $\iota = \mathrm{id}_V$ and $f^0 = \iota$. Let $\langle x\rangle_f = \mathrm{span}\{f^i x, i \geq 0\}$ be the cyclic subspace generated by $x \in V$. If $B \subseteq V$ we define $\langle B\rangle_f = \sum_{b\in B} \langle\, b\,\rangle_f$. Let λ be an eigenvalue of f such that $V_\lambda = \mathrm{Ker}(f - \lambda\iota)^n$ is the corresponding generalized eigenspace. Let $\dim \mathrm{Ker}(f - \lambda\iota) = k$, and let $s^{t_1}, \ldots, s^{t_k}$, be the elementary divisors of $f_{|V_\lambda}$. Then there exist vectors $u_1, \ldots, u_k$, such that

$$V_\lambda = \langle u_1\rangle_{f-\lambda\iota} \oplus \cdots \oplus \langle u_k\rangle_{f-\lambda\iota},$$

and $(f - \lambda\iota)^{t_i-1}u_i \neq 0$, $(f - \lambda\iota)^{t_i}u_i = 0$, $i = 1, \ldots, k$. We call $U_\lambda = \{u_1, \ldots, u_k\}$ a set of *generators* of V_λ. Each U_λ gives rise to a Jordan basis of V_λ, namely

$$\big\{u_1,\, (f - \lambda\iota)u_1,\, \ldots, (f - \lambda\iota)^{t_1-1}u_1 \ldots,\; u_k,\, (f - \lambda\iota)u_k,\, \ldots, (f - \lambda\iota)^{t_k-1}u_k\big\}.$$

Define $f_\lambda = f_{|V_\lambda}$. Let Y be an f_λ-invariant subspace of V_λ. Then Y is said to be *marked* in V_λ (with respect to f_λ) if there exists a set U_λ of generators of V_λ and corresponding integers r_i, $0 \leq r_i \leq t_i$, such that

$$Y = \big\langle (f - \lambda\iota)^{r_1} u_1\big\rangle_{f-\lambda\iota} \oplus \cdots \oplus \big\langle (f - \lambda\iota)^{r_k} u_k\big\rangle_{f-\lambda\iota}.$$

Thus Y has a Jordan basis which can be extended to a Jordan basis of V_λ. Let $\sigma(f) = \{\lambda_1, \ldots, \lambda_m\}$ be the spectrum of f. Then

$$V = V_{\lambda_1} \oplus \cdots \oplus V_{\lambda_m}. \tag{1.1}$$

If $X \in \mathrm{Inv}_f V$ then $X_{\lambda_i} = X \cap V_{\lambda_i}$ is f_{λ_i}-invariant in V_{λ_i}, and

$$X = X_{\lambda_1} \oplus \cdots \oplus X_{\lambda_m}. \tag{1.2}$$

We say that X is marked in V if each subspace X_{λ_i} in (1.2) is marked in V_{λ_i}. The set of marked subspaces of V will be denoted by Mark(V). We assume $0 \in$ Mark(V). Marked subspaces can be traced back to [13, p. 83]. They have been studied in [4], [8], [1], and [6]. For marked (A, C)-invariant subspaces we refer to [5] and [7]. We mention applications to algebraic Riccati equations [2] and to stability of invariant subspaces of commuting matrices [15].

The following examples show that to a certain extent the three types of invariant subspaces are independent of each other. Suppose f is nilpotent. If $x \in V$ then the smallest nonnegative integer ℓ with $f^\ell x = 0$ is called the *exponent* of x. We write $\mathrm{e}(x) = \ell$. A nonzero vector x is said to have *height* q if $x \in f^q V$ and $x \notin f^{q+1}V$. In this case we write $\mathrm{h}(x) = q$. We set $\mathrm{h}(0) = -\infty$. For $j \geq 0$ we define $V[f^j] = \mathrm{Ker}\, f^j$.

Example 1.1. Let $K = \mathbb{Z}_2$. Consider $V = K^4$ and

$$f = \mathrm{diag}(0, N_3), \;\; N_3 = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}.$$

Let $e_1, \ldots, e_4$, be the unit vectors of K^4. Then $f^3 = 0$ and $V = \langle e_1 \rangle_f \oplus \langle e_2 \rangle_f$. Define $z = e_1 + e_3$ and $Z = \langle z \rangle_f$. Then

$$Z = \{0, z, z + e_4, e_4\} = \big\langle v;\ \mathrm{e}(v) = 2,\ \mathrm{h}(v) = 0,\ \mathrm{h}(fv) = 2 \big\rangle_f.$$

If $\alpha \in \mathrm{Aut}_f(V)$ then $|\alpha(Z)| = |Z|$. Moreover α preserves height and exponent. Hence $\alpha(Z) = Z$, and Z is characteristic. Let $g = \mathrm{diag}(1, 0, 0, 0)$ be the orthogonal projection on Ke_1. Then $g \in \mathrm{End}_f(V)$. We have $gz = e_1 \in g(Z)$, but $e_1 \notin Z$. Therefore Z is not hyperinvariant. The Jordan bases of Z are $J_1 = \{z, e_4\}$ and $J_2 = \{z + e_4, e_4\}$. If $y \in K^4$ then $z \neq fy$ and $z + e_4 \neq fy$. Hence neither J_1 nor J_2 can be extended to a Jordan basis of K^4. Therefore Z is not marked.

Example 1.2. Let $V = K^2$ and $f = 0$. Then $K^2 = \langle e_1 \rangle_f \oplus \langle e_2 \rangle_f$ and the subspace $X = \langle e_1 \rangle_f$ is marked. From $\alpha = \left(\begin{smallmatrix} 1 & 0 \\ 1 & 1 \end{smallmatrix}\right) \in \mathrm{Aut}_f(V)$ and $\alpha(e_1) = e_1 + e_2$ follows that X is not characteristic.

In contrast to Hinv(V) or Chinv(V) the set Mark(V) in general is not a lattice.

Example 1.3. $V = K^6$, $f = \mathrm{diag}(0, N_3, N_2)$. The subspaces $Z_1 = \langle e_5 \rangle$ and $Z_2 = \langle e_5 + e_3 + e_1 \rangle$ are marked but $Z_1 + Z_2 = \langle e_5 \rangle \oplus \langle e_3 + e_1 \rangle$ is not marked. Thus the set of marked subspaces is not closed under addition.

In this paper we study the following problems. Under what conditions is a marked subspace characteristic? When is each characteristic subspace hyperinvariant? Because of the Lemma 1.4 below one can deal separately with single components V_{λ_i} in (1.1) and the corresponding restrictions $f_{\lambda_i} = f_{|V_{\lambda_i}}$, $i = 1, \ldots, m$.

Lemma 1.4. *An f-invariant subspace $X \subseteq V$ is hyperinvariant (resp. characteristic, resp. marked) if and only if, with respect to f_{λ_i}, each component X_{λ_i} in* (1.2) *is hyperinvariant (resp. characteristic, resp. marked) in V_{λ_i}.*

Proof. If $\eta \in \mathrm{End}_f(V)$ then it is known ([11, p. 223]) that the subspaces V_{λ_i} in (1.1) are invariant under η, and that $\eta_{|V_{\lambda_i}} \in \mathrm{End}_{f_{\lambda_i}}(V_{\lambda_i})$. Hence, if $X \in \mathrm{Inv}(V)$ then (1.2) implies

$$\eta(X) = \eta_{|V_{\lambda_1}}(X_{\lambda_1}) \oplus \cdots \oplus \eta_{|V_{\lambda_i}}(X_{\lambda_m})$$

Hence if $X_{\lambda_i} \in \mathrm{Hinv}(V_{\lambda_i})$, resp. $X_{\lambda_i} \in \mathrm{Chinv}(V_{\lambda_i})$, $i = 1, \ldots, m$, then $X \in \mathrm{Hinv}(V)$, resp. $X \in \mathrm{Chinv}(V)$.

Now suppose now that X is hyperinvariant. Let us show that $X_{\lambda_i} \in \mathrm{Hinv}(V_{\lambda_i})$, $i = 1, \ldots, m$. Take $i = 1$. Set $\hat{V} = V_{\lambda_2} \oplus \cdots \oplus V_{\lambda_m}$ and $\hat{X} = X_{\lambda_2} \oplus \cdots \oplus X_{\lambda_m}$. Let $\beta_1 \in \mathrm{End}_{f_{\lambda_1}}(V_{\lambda_1})$. Define $\beta = \beta_1 + \mathrm{id}_{|\hat{V}}$. Then $\beta \in \mathrm{End}_f(V)$. Hence $\beta(X) \subseteq X = X_{\lambda_1} \oplus \hat{X}$, and $\beta(X) = \beta_1(X_{\lambda_1}) \oplus \hat{X}$. From $X_{\lambda_1} \subseteq V_{\lambda_1}$ and $\beta_1(X_{\lambda_1}) \subseteq V_{\lambda_1}$ we obtain $\beta_1(X_{\lambda_1}) \subseteq X_{\lambda_1}$. Therefore $X_{\lambda_1} \in \mathrm{Hinv}(V_{\lambda_1})$. A similar argument shows that $X \in \mathrm{Chinv}(V)$ implies $X_{\lambda_i} \in \mathrm{Chinv}(V_{\lambda_i})$, $i = 1, \ldots, m$. In the case of marked subspaces the assertion is obvious. □

2 Auxiliary results

Because of Lemma 1.4 it suffices to consider an endomorphism f with only one eigenvalue λ. We shall assume $\sigma(f) = \{0\}$ such that $f^n = 0$. Let

$$s^{t_1}, \ldots, s^{t_k}, \quad 0 < t_1 \leq \cdots \leq t_k, \tag{2.1}$$

be the elementary divisors of f. We call $U = (u_1, \ldots, u_k)$ a *generator tuple* of V if

$$V = \langle u_1 \rangle_f \oplus \cdots \oplus \langle u_k \rangle_f \tag{2.2}$$

and if U is ordered according to (2.1) such that

$$\mathrm{e}(u_1) \;=\; t_1 \;\leq\; \cdots \;\leq\; \mathrm{e}(u_k) \;=\; t_k.$$

Let $\mathcal{U}$ be the set of generator tuples of V. In the following we omit the subscript f in (2.2) and we write $\langle u_i \rangle$ instead of $\langle u_i \rangle_f$. We say that a k-tuple $r = (r_1, \ldots, r_k)$ of integers is *admissible* if

$$0 \leq r_i \leq t_i, \;\; i = 1, \ldots, k. \tag{2.3}$$

Each $U \in \mathcal{U}$ together with an admissible tuple r gives rise to a subspace

$$W(r, U) = \langle f^{r_1} u_1 \rangle \;\oplus\; \cdots \;\oplus\; \langle f^{r_k} u_k \rangle, \tag{2.4}$$

which is marked in V. Conversely, a subspace W is marked in V only if $W = W(r, U)$ for some $U \in \mathcal{U}$ and some admissible r. The following example shows that, in general, $W(r, U) \neq W(r, \tilde{U})$ if $U \neq \tilde{U}$.

Example 2.1. Let $V = K^5$ and $f = \mathrm{diag}(N_2, N_3)$. Then $V = \langle e_1 \rangle \oplus \langle e_3 \rangle$, and $U = (e_1, e_3)$ and $\tilde{U} = (e_1, e_3 + e_1)$ are generator tuples. Choose $r = (1, 0)$. Then the corresponding subspaces $W(r, U) = \langle e_2 \rangle \oplus \langle e_3 \rangle$ and $W(r, \tilde{U}) = \langle e_2 \rangle \oplus \langle e_3 + e_1 \rangle$ are different from each other.

The construction of invariant subspaces of the form $W(r, U)$ is a standard procedure in linear algebra and systems theory. It is used in [16], [12, p.61], [3, p.28], [18]. Hence it is important to know whether for a given r different choices of U will always result in the same subspace. Theorem 3.1 will provide a necessary and sufficient condition for r such that $W(r, U)$ is independent of the choice of U. Let r be admissible and define

$$W(r) = f^{r_1} V \cap V[f^{t_1 - r_1}] \;+ \cdots + \; f^{r_k} V \cap V[f^{t_k - r_k}]. \tag{2.5}$$

Subspaces of the form $f^\nu V$ and $V[f^\mu]$ are hyperinvariant, and $\mathrm{Hinv}(V)$ is a lattice. Therefore (see e.g. [9]) we have $W(r) \in \mathrm{Hinv}(V)$.

The following lemma shows that each $\alpha \in \mathrm{Aut}_f(V)$ is uniquely determined by the image of a given generator tuple.

Lemma 2.2. *Let $U = (u_1, \ldots, u_k) \in \mathcal{U}$ be given. For $\alpha \in \mathrm{Aut}_f(V)$ define $\Theta_U(\alpha) = \big(\alpha(u_1), \ldots, \alpha(u_k)\big)$.* (i) *Then*

$$\alpha \mapsto \Theta_U(\alpha), \;\; \Theta_U : \mathrm{Aut}_f(V) \to \mathcal{U},$$

is a bijection. (ii) *If $\tilde{U} = \Theta(\alpha)$ then $W(r, \tilde{U}) = \alpha\big(W(r, U)\big)$.*

Proof. (i) It is easy to see that $\Theta_U(\alpha) \in \mathcal{U}$. Hence Θ_U maps $\mathrm{Aut}_f(V)$ into $\mathcal{U}$. Let $x \in V$ and

$$x = \sum\nolimits_{i=1}^{k} \sum\nolimits_{j=0}^{\mathrm{e}(u_i)-1} c_{ij} f^j u_i. \tag{2.6}$$

Suppose $\alpha, \beta \in \mathrm{Aut}_f(V)$ and $\Theta_U(\alpha) = \Theta_U(\beta) = (\hat{u}_1, \ldots, \hat{u}_k)$. Then

$$\alpha(x) = \sum \sum c_{ij} f^j \hat{u}_i = \beta(x).$$

Hence $\alpha = \beta$, and Θ_U is injective. Now consider $\tilde{U} = (\tilde{u}_1, \ldots, \tilde{u}_k) \in \mathcal{U}$. Let $x \in V$ be the vector in (2.6). Define $\gamma : x \mapsto \sum_i \sum_j c_{ij} f^j \tilde{u}_i$. Then $\gamma \in \mathrm{Aut}_f(V)$ and $\tilde{U} = \Theta_U(\gamma)$. Hence Θ_U is surjective.
(ii) It is obvious that $\alpha\big(W(r,U)\big) = \langle f^{r_1}\alpha(u_1)\rangle_f \oplus \cdots \oplus \langle f^{r_k}\alpha(u_k)\rangle_f = W(r,\tilde{U})$. □

In group theory fully invariant subgroups play the role of hyperinvariant subspaces. Hence the decomposition (2.8) below is an analog to a distributive law in Lemma 9.3 in [10, p. 47].

Lemma 2.3. *Suppose*

$$V = V_1 \oplus \cdots \oplus V_q,\ V_i \in \mathrm{Inv}(V),\ i = 1, \ldots, q. \tag{2.7}$$

(i) *If X is a hyperinvariant subspace of V, or*
(ii) *if X characteristic and $|K| > 2$, then*

$$X = (X \cap V_1) \oplus \cdots \oplus (X \cap V_q). \tag{2.8}$$

Proof. If $x \in V$ then $x = \sum_{i=1}^{q} x_i$, $x_i \in V_i$. Set $X_i = X \cap V_i$, and $S = \oplus_{i=1}^{q} X_i$. Then $S \subseteq X$. To prove the converse inclusion we note that

$$fx = \sum\nolimits_{i=1}^{q} f_{|V_i}(x_i). \tag{2.9}$$

(i) Let π_i be the projection on V_i induced by (2.7). Then (2.9) implies $\pi_i \in \mathrm{End}_f(V)$. Hence, if $x \in X$ then and $\pi_i(x) = x_i \in X$. Thus $x_i \in X_i$, and therefore $X \subseteq S$.
(ii) Let $a \in K$ be different from 0 and 1, and define $\gamma_i = \iota - a\pi_i$. Then $\gamma_i \in \mathrm{Aut}_f(V)$. Hence $\gamma_i(x) = x - ax_i \in X$ if $x \in X$. Thus we obtain $x_i \in X_i$. □

Example 2.4. In Lemma 2.3(ii) one can not drop the assumption $|K| > 2$. Suppose $|K| = 2$, and let V and f be as in Example 1.1. The subspace $Z = \langle e_1 + e_3 \rangle$ is characteristic. Both $V_1 = \langle e_1 \rangle$ and $V_2 = \langle e_2 \rangle$ are in $\mathrm{Inv}(V)$, and we have $V = V_1 \oplus V_2$. But $Z \cap V_1 = 0$ and $Z \cap V_2 = \langle e_4 \rangle$ imply $Z \supsetneqq (Z \cap V_1) \oplus (Z \cap V_2)$.

The next lemma is an intermediate result.

Lemma 2.5. *Each hyperinvariant subspace of V is marked, and*

$$\mathrm{Hinv}(V) \subseteq \mathrm{Mark}(V) \cap \mathrm{Chin}(V). \tag{2.10}$$

Proof. Let $U = (u_1, \ldots, u_k) \in \mathcal{U}$. If X is invariant then $X \cap \langle u_i \rangle = \langle f^{r_i} u_i \rangle$ for some r_i. Thus, if X is hyperinvariant then (2.8) in Lemma 2.3 implies $X = \oplus_{i=1}^{k} \langle f^{r_i} u_i \rangle$. Therefore X is marked, and $\mathrm{Hinv}(V) \subseteq \mathrm{Chin}(V)$ yields the inclusion (2.10). □

3 Hyperinvariant = characteristic + marked

We now characterize those marked subspaces which are characteristic. The theorem below includes results from [2] with new proofs.

Theorem 3.1. *Let $U \in \mathcal{U}$ and let $r = (r_1, \ldots, r_k)$ be admissible. Then the following statements are equivalent.*

(i) *The subspace $W(r, U)$ is characteristic.*

(ii) *The subspace $W(r, U)$ is independent of the generator tuple U, i.e.*

$$W(r, U) = W(r, \tilde{U}) \quad \textit{for all} \quad \tilde{U} \in \mathcal{U}. \tag{3.1}$$

(iii) *The tuples $t = (t_1, \ldots, t_k)$ and $r = (r_1, \ldots, r_k)$ satisfy*

$$r_1 \leq \cdots \leq r_k \tag{3.2}$$

and

$$t_1 - r_1 \leq \cdots \leq t_k - r_k. \tag{3.3}$$

(iv) *We have $W(r, U) = W(r)$.*

(v) *$W(r, U)$ is the unique marked subspace W such that the elementary divisors of W and of V/W are*

$$s^{t_1 - r_1}, \ldots, s^{t_k - r_k}, \quad \text{and} \quad s^{r_1}, \ldots, s^{r_k}. \tag{3.4}$$

(vi) *The subspace $W(r, U)$ is hyperinvariant.*

Proof. (i) $\Leftrightarrow$ (ii) It follows from Lemma 2.2 that the two statements are equivalent.
(iv) $\Rightarrow$ (vi) This follows from the fact that $W(r)$ is hyperinvariant.
(v) $\Leftrightarrow$ (ii) Let $\tilde{U} \in \mathcal{U}$. Then $W(r,U)$ and the quotient space $V/W(r,U)$, and also $W(r,\tilde{U})$ and $V/W(r,\tilde{U})$, have elementary divisors given by (3.4). (Note that in the right-hand side of (2.4) there may be summands of the form $\langle u_i\rangle$ or $\langle f^{t_i}u_i\rangle = 0$. Thus (3.4) may contain trivial entries of the form $s^0 = 1$.)
(vi) $\Rightarrow$ (i) Obvious, because of $\mathrm{Hinv(V)} \subseteq \mathrm{Chinv(V)}$.
(iii) $\Rightarrow$ (iv) From $\mathrm{e}(u_i) = t_i$ follows

$$\langle f^{r_i}u_i\rangle = \langle u_i\rangle[f^{t_i-r_i}] \subseteq f^{r_i}V \cap V[f^{t_i-r_i}].$$

Hence $W(r,U) \subseteq W(r)$. We have to show that the conditions (3.2) and (3.3) imply the converse inclusion

$$W(r) = f^{r_1}V \cap V[f^{t_1-r_1}] + \cdots + f^{r_k}V \cap V[f^{t_k-r_k}] \subseteq W(r,U).$$

With regard to the decomposition $V = \langle u_1\rangle \oplus \cdots \oplus \langle u_k\rangle$ we define

$$D(\mu,\nu) = f^{r_\nu}\langle u_\mu\rangle \cap \langle u_\mu\rangle\,[f^{t_\nu-r_\nu}].$$

The subspaces $f^{r_\nu}V \cap V[f^{t_\nu-r_\nu}]$ are hyperinvariant. Therefore Lemma 2.3(i) yields

$$f^{r_\nu}V \cap V[f^{t_\nu-r_\nu}] = \bigoplus_{\mu=1}^{k}\big(f^{r_\nu}V \cap V[f^{t_\nu-r_\nu}] \cap \langle u_\mu\rangle\big) = \bigoplus_{\mu=1}^{k} D(\mu,\nu).$$

Hence

$$W(r) = \sum_{\mu,\nu=1}^{k} D(\mu,\nu). \tag{3.5}$$

Set $q(\mu,\nu) = \max\{r_\nu,\, t_\mu - (t_\nu - r_\nu)\}$. We have

$$\langle u_\mu\rangle[f^{t_\nu-r_\nu}] = \begin{cases} \langle u_\mu\rangle, & \text{if } t_\nu - r_\nu \geq t_\mu, \\ f^{t_\mu-(t_\nu-r_\nu)}\langle u_\mu\rangle, & \text{if } t_\nu - r_\nu \leq t_\mu. \end{cases}$$

Hence

$$D(\mu,\nu) = f^{q(\mu,\nu)}\langle u_\mu\rangle.$$

Let us show that $r_\mu \leq q(\mu,\nu)$ for all μ. If $\mu \geq \nu$, then (3.3) implies

$$q(\mu,\nu) = (t_\mu - t_\nu) + r_\nu = (t_\mu - r_\mu) - (t_\nu - r_\nu) + r_\mu \geq r_\mu.$$

If $\mu \leq \nu$ then $t_\mu - t_\nu \leq 0$, and therefore $q(\mu,\nu) = r_\nu$. Hence (3.2) implies $q(\mu,\nu) \geq r_\mu$. It follows that

$$D(\mu,\nu) = f^{q(\mu,\nu)}\langle u_\mu\rangle \subseteq f^{r_\nu}\langle u_\mu\rangle \subseteq W(r,U).$$

for all μ,ν. Thus (3.5) yields $W(r) \subseteq W(r,U)$.
(ii) $\Rightarrow$ (iii) We modify the entries of U and replace u_k by $\tilde{u}_k = u_{k-1}+u_k$. Then $\tilde{U} = (u_1,\ldots,u_{k-1},\tilde{u}_k) \in \mathcal{U}$. Set $Y_k = \oplus_{i=1}^{k-1}\langle f^{r_i}u_i\rangle$. Then $W(r,U) = W(r,\tilde{U})$ implies

$$Y_k \oplus \langle f^{r_k}u_k\rangle = Y_k \oplus \langle f^{r_k}(u_{k-1}+u_k)\rangle.$$

From

$$f^{r_k}u_{k-1} + f^{r_k}u_k \ \in \ \langle f^{r_{k-1}}u_{k-1}\rangle \oplus \langle f^{r_k}u_k\rangle$$

follows $r_{k-1} \leq r_k$. Proceeding in this manner we obtain the chain of inequalities in (3.2). In order to prove (3.3) we start with the entry of u_1 of $\mathcal{U}$ and replace it by $u_1 + f^{t_2-t_1}u_2$. Because of $\mathrm{e}(u_1 + f^{t_2-t_1}u_2) = \mathrm{e}(u_1)$ we have $\hat{U} = (u_1 + f^{t_2-t_1}u_2, u_2, \ldots, u_k) \in \mathcal{U}$. Set $Y_1 = \oplus_{i=2}^{k}\langle f^{r_i}u_i\rangle$. Then $W(r,U) = W(r,\hat{U})$ implies

$$\langle f^{r_1}u_1\rangle \oplus Y_1 = \langle f^{r_1}(u_1 + f^{t_2-t_1}u_2)\rangle \oplus Y_1.$$

From

$$f^{r_1}u_1 + f^{r_1+(t_2-t_1)}u_2 \ \in \ \langle f^{r_1}u_1\rangle \oplus \langle f^{r_2}u_2\rangle$$

follows $r_2 \leq r_1 + (t_2 - t_1)$, i.e. $t_1 - r_1 \leq t_2 - r_2$, such that we the end up with (3.3). □

Let $[k]$ denote the greatest integer less than or equal to k. If $c \in \mathbb{R}$ and $0 < c < 1$, then $r = ([c\,t_1],\ldots,[c\,t_m])$ is admissible, and it is not difficult to verify that r satisfies (3.2) and (3.3). We remark that admissible tuples of the form $\hat{r} = ([\frac{1}{2}t_1],\ldots,[\frac{1}{2}t_k])$ play a role in the study of maximal invariant neutral subspaces [18]. It follows from Theorem 3.1 that the construction of such subspaces is independent of the choice of the underlying Jordan basis.

Theorem 3.2. (i) *We have*

$$\mathrm{Hinv}(V) = \mathrm{Chinv}(V) \cap \mathrm{Mark}(V). \tag{3.6}$$

(ii) [9] *A subspace W of V is hyperinvariant if and only if $W = W(r)$ for some r satisfying* (3.2) *and* (3.3).

Proof. (i) From Theorem 3.1 follows $\mathrm{Mark}(V) \cap \mathrm{Chinv}(V) \subseteq \mathrm{Hinv}(V)$. The reverse inclusion is (2.10) in Lemma 2.5. This yields (3.6). Hence a subspace is hyperinvariant if and only if it is both characteristic and marked.
(ii) If W is hyperinvariant then W is marked, that is $W = W(r, U)$. Therefore we can apply Theorem 3.1(iv). It was noted earlier that $W(r) \in \mathrm{Hinv}(V)$. □

We note that hyperinvariant subspaces can be characterized completely by the distributive law in Lemma 2.3.

Theorem 3.3. *A subspace $X \in \mathrm{Inv}(V)$ is hyperinvariant if and only if X satisfies*

$$X = (X \cap V_1) \oplus \cdots \oplus (X \cap V_q) \tag{3.7}$$

when

$$V = V_1 \oplus \cdots \oplus V_q,\ V_i \in \mathrm{Inv}(V),\ i = 1, \ldots, q. \tag{3.8}$$

Proof. Because of Lemma 2.3 it remains to prove sufficiency. Let $U = (u_1, \ldots, u_k) \in \mathcal{U}$ and $\tilde{U} = (\tilde{u}_1, \ldots, \tilde{u}_k) \in \mathcal{U}$. Then

$$V = \langle u_1 \rangle \oplus \cdots \oplus \langle u_k \rangle = \langle \tilde{u}_1 \rangle \oplus \cdots \oplus \langle \tilde{u}_k \rangle. \tag{3.9}$$

Define $X_i = \langle u_i \rangle \cap X$ and $\tilde{X}_i = \langle \tilde{u}_i \rangle \cap X$, $i = 1, \ldots, k$. Then $X_i = \langle f^{r_i} u_i \rangle$ and $\tilde{X}_i = \langle f^{\tilde{r}_i} \tilde{u}_i \rangle$ for some $r_i, \tilde{r}_i$. Set $r = (r_1, \ldots, r_k)$ and $\tilde{r} = (\tilde{r}_1, \ldots, \tilde{r}_k)$. In (3.9) we have two direct sums of the form (3.8). Hence the assumption (3.7) implies $X = W(r, U) = W(\tilde{r}, \tilde{U})$. We can pass from U to $\tilde{U}$ in at most k steps, changing a single entry at each step. Suppose we replace u_k in U by $\tilde{u}_k$. Then $\hat{U} = (u_1, \ldots, u_{k-1}, \tilde{u}_k) \in \mathcal{U}$, and $V = \langle u_1 \rangle \oplus \cdots \langle u_{k-1} \rangle \oplus \langle \tilde{u}_k \rangle$. Set $Y_k = \oplus_{i=1}^{k-1} \langle f^{r_i} u_i \rangle$. Then

$$X = Y_k \oplus \langle f^{\tilde{r}_k} \tilde{u}_k \rangle = Y_k \oplus \langle f^{r_k} u_k \rangle.$$

Considering the elementary divisors of V/X we deduce $\tilde{r}_k = r_k$, and at the end we obtain $r = \tilde{r}$, and therefore $W(r, U) = W(r, \tilde{U})$. We conclude that $X = W(r, U)$ is independent of the choice of the generator tuple U. Hence X is hyperinvariant. □

Let us reexamine Example 1.1 and consider a field K of characteristic different from 2.

Example 1.1 (continued). Let $\mathrm{char} K \neq 2$. Then $\gamma : (e_1, e_2) \mapsto (2e_1, e_2)$ determines an f-automorphism. For $Z = \langle e_1 + e_3 \rangle$ we have $\gamma(Z) = \langle 2e_1 + e_3 \rangle \neq Z$. Hence in this case $Z \in \mathrm{Inv}(V)$ is not characteristic.

To identify the characteristic subspaces we screen $\mathrm{Inv}(V)$. Note that

$$\mathrm{Aut}_f(V) = \big\{\alpha : (e_1, e_2) \mapsto (ae_1 + be_4, ce_2 + de_3 + ge_4 + he_1), \\ a, b, c, d, g, h \in K, a \neq 0, c \neq 0\big\}.$$

The nonzero cyclic subspaces are of the form $\langle e_2 + ce_1\rangle$, $\langle e_3 + ce_1\rangle$, and $\langle ae_4 + ce_1\rangle$, $a, c \in K$, $(a, c) \neq (0, 0)$. Only $\langle e_3\rangle = fV$ and $\langle e_4\rangle = f^2V$ are characteristic. Moreover, X is a direct sum of two cyclic subspaces if and only if $X \in \{V, \langle e_3\rangle \oplus \langle e_1\rangle = V[f^2], \langle e_4\rangle \oplus \langle e_1\rangle = V[f]\}$. These three subspaces are characteristic. We find $\mathrm{Hinv}(V) = \{0,\ fV,\ f^2V,\ V[f],\ V[f^2],\ V\}$. Hence $\mathrm{Hinv}(V) = \mathrm{Chinv}(V)$. The example is a special case of the following general result (see also [14, p. 67]).

Theorem 3.4. *If* $|K| > 2$ *then each characteristic subspace of* V *is hyperinvariant, i.e.* $\mathrm{Chinv}(V) = \mathrm{Hinv}(V)$.

Proof. Because of Lemma 1.4 it suffices to consider the case where f has only one eigenvalue. We can assume $f^n = 0$. If $|K| > 2$ and X is characteristic then it follows from Lemma 2.3(ii) that (2.7) implies (2.8). Therefore, according to Theorem 3.3, the subspace X is hyperinvariant. □

In the case of vector spaces over $K = \mathbb{Z}_2$ it is an open problem to describe all subspaces that are characteristic without being hyperinvariant.

Acknowledgment: We are indebted to L. Rodman for a valuable remark.

References

[1] P. Astuti and H. K. Wimmer, Regular submodules of torsion modules over a discrete valuation domain, Czechoslov. Math. J. 56(2006), 349–357.

[2] P. Astuti and H. K. Wimmer, A class of marked invariant subspaces with an application to algebraic Riccati equations, Automatica 42 (2006), 1503–1506.

[3] L. Baratchart, P. Enqvist, A. Gombani, and M. Olivi, Surface acoustic wave filters, unitary extensions and Schur analysis, Report No. 44, 2003, Institut Mittag-Leffler, Stockholm, http://www.mittag-leffler.se/preprints/0203s.

[4] R. Bru, L. Rodman, and H. Schneider, Extensions of Jordan bases for invariant subspaces of a matrix, Linear Algebra Appl. 150(1991), 209–226.

[5] A. Compta and J. Ferrer, On $(A, B)^t$-invariant subspaces having extendible Brunovsky bases, Linear Algebra Appl. 255(1997), 185–201.

[6] A. Compta, J. Ferrer, and M. Peña, Dimensions of the orbit of marked subspaces, Linear Algebra Appl. 379(2004), 239–248.

[7] A. Compta, and M. Peña, Structural stability of (C, A)-marked subspaces, Linear Algebra Appl. 421(2007), 45–52.

[8] J. Ferrer, F. Puerta, and X. Puerta, Geometric characterization and classification of marked subspaces, Linear Algebra Appl. 235(1996), 15–34.

[9] P. A. Fillmore, D. A. Herrero, and W. E. Longstaff, The characteristic subspace lattice of a linear transformation, Linear Algebra Appl. 17(1977), 125–132.

[10] L. Fuchs, Infinite Abelian Groups, Vol. I., Academic Press, New York, 1973.

[11] F. R. Gantmacher, The Theory of Matrices, Vol. I, Chelsea, New York 1959.

[12] I. Gohberg, P. Lancaster, and L. Rodman, Matrices and Indefinite Scalar Products, Operator Theory: Advances and Applications, Vol. 8, Birkhäuser, Basel, 1983.

[13] I. Gohberg, P. Lancaster, and L. Rodman, Invariant Subspaces of Matrices with Applications, Wiley, New York, 1986.

[14] I. Kaplansky, Infinite Abelian Groups, University of Michigan Press, Ann Arbor, 1954.

[15] T. Košir and B. Plestenjak, On stability of invariant subspaces of commuting matrices, Linear Algebra Appl. 342(2002), 133–147.

[16] V. Kučera, Algebraic Riccati equations: Hermitian and definite solutions, in S. Bittanti, A.J. Laub and J.C. Willems (Eds.), *The Riccati Equation*, pp.53–88, Springer Verlag, Berlin 1991.

[17] W. E. Longstaff, A lattice-theoretic description of the lattice of hyperinvariant subspaces of a linear transformation, Can. J. Math. 28(1976), 1062–1066.

[18] L. Rodman, Maximal invariant neutral subspaces and an application to the algebraic Riccati equation, Manuscr. Math. 43(1983), 1–12.

[19] P. Y. Wu, Which linear transformations have isometric hyperinvariant subspace lattices?, Linear Algebra Appl. 169(1992), 163–178.